


# Studies of Particle Formation, Growth, and Chemistry Using Aerosol Mass Spectrometry

Jesse Kroll

Depts. of Civil and Environmental Engineering and Chemical Engineering  
Massachusetts Institute of Technology

Workshop on Techniques for High-Pressure Combustion

31 August 2011



# Atmospheric chemistry

Conditions:  $P \leq 1$  bar,  $T \leq 320$  K,  $\chi \leq$  ppb, ppt

Relation to combustion chemistry:

- 1) Emissions
- 2) Similarities in terms of (organic) chemistry
  - Radical-initiated oxidation of species to form low-volatility (particulate) material
  - Multiphase processes: homogeneous reactions, heterogeneous reactions, partitioning between phases
  - Many reactants, intermediates, products → immensely complex chemical mechanisms

# Outline


---

- 1) Atmospheric aerosol, aerosol mass spectrometry
- 2) Flow tube studies of aerosol chemistry
- 3) Carbon oxidation state of organics
- 4) Connection with combustion chemistry

# Atmospheric aerosol


Fine particles: diameter  $\leq 1\mu\text{m}$

## 1) Human health



D. W. Dockery et al., *New Engl. J. Med.* 329, 1753-1759 (1993)


## 2) Global climate




IPCC: *Climate Change 2007: The Physical Science Basis* (2007).

# Atmospheric aerosol loadings/composition

Fine particles ("aerosols"):  $d_p \leq 1 \mu\text{m}$




# Aerodyne Aerosol Mass Spectrometer (AMS)




Quantitative, real-time measurement of aerosol mass, composition

- (1) Particle focusing/concentration
- (2) Particle sizing
- (3) Vaporization at 600°C, electron impact ionization at 70 eV (ion fragmentation)
- (4) High-resolution (5000) time-of-flight mass spectrometry

# High-resolution electron impact AMS




# Vacuum ultraviolet photoionization (VUV) AMS



VUV from the chemical dynamics beamline (9.0.2) at the Advanced Light Source,  
Lawrence Berkeley National Laboratory [Kevin Wilson, Steve Leone, et al.]


“Soft ionization” approach: vaporizer at <100°C, photon energy (8-12 eV) just  
above ionization threshold (~9-10 eV for large organics)

# AMS: Hard (EI) vs. soft (VUV) ionization




# Flow tube experiments (heterogeneous oxidation)

Flow reactor at Lawrence Berkeley National Laboratory (PI: Kevin Wilson):





- AMS + scanning mobility particle sizer (SMPS): elemental ratios, particle volume, density, mass
- $[\text{OH}]$  varied by changing  $[\text{O}_3]$ : can access hours to weeks of atmospheric oxidation


# VUV-AMS spectrum: Oxidized squalane





# VUV-AMS spectrum: Oxidized squalane



+1O, -2H





# VUV-AMS spectrum: Oxidized squalane




Some fraction of the aerosol becomes highly oxygenated (>5 O atoms) very quickly


# Sequential oxidation



$$\frac{[\text{SqO}_n]}{[\text{Sq}]_0} = B_n \frac{(k \cdot \langle \text{OH} \rangle \cdot t)^n}{n!} \text{Exp}(-k \cdot \langle \text{OH} \rangle_t \cdot t)$$




# Ensemble measurements: Physical changes




density increases  
volume decreases  
→ net decrease in mass

# Ensemble measurements: Chemical changes




O/C increases  
H/C decreases  
→ oxidation state increases

# Elemental abundances




# Oxidation of organic species

In an oxidizing environment, the oxidation state of carbon increases:



# Carbon oxidation state vs. carbon number



# Measuring carbon oxidation state

Trivial for speciated measurements

Can also be determined for complex mixtures:

$$\overline{OS}_C = -\sum_i OS_i \frac{n_i}{n_C}$$


$OS_i$  = oxidation state of atom  $i$

$n_i$  = number of atom  $i$

Organic aerosol is made up largely of carbon, hydrogen ( $OS=+1$ ) and reduced oxygen ( $OS=-2$ ) :


$$\overline{OS}_C \approx 2 \times O/C - H/C$$

# Oxidation state evolution



squalane  $C_{30}H_{62}$

# Oxidation state evolution





squalane  $C_{30}H_{62}$



triacontane  $C_{30}H_{62}$

# Oxidation state evolution



squalane  $C_{30}H_{62}$



triacontane  $C_{30}H_{62}$






coronene  $C_{24}H_{12}$   
+ OH



coronene  $C_{24}H_{12}$   
+  $O_3$


# Oxidation state evolution




+ OH



+  $O_3$




# Combustion processes



# AMS measurements of soot in combustion systems

Challenges:

- 1) Interface: high P, T
- 2) Refractory nature of soot: won't flash vaporize at 600°C
- 3) Very small particles (<10 nm): can't be focused in aerodynamic lens



# Conclusions

Challenges in studying the chemistry of organic aerosol (OA) include:

- multistep, multiphase nature of the chemical mechanisms
- immense chemical complexity

We can understand key features of complex mechanisms and condensed-phase mixtures using *highly time-resolved aerosol mass spectrometry*:

- hard ionization (EI): ensemble measurements
- soft ionization (VUV): speciated measurements

Carbon oxidation state ( $OS_C$ ) and carbon number ( $n_C$ ) serve as a useful ‘reduced representation’ for describing atmospheric OA chemistry

Connections between atmospheric chemistry and combustion chemistry (and especially between OA and soot) → potential opportunities for both communities


# Acknowledgements



Sean Kessler  
Kelly Daumit



**Kevin Wilson**  
Jared Smith  
Theodora Nah



Doug Worsnop  
Tim Onasch  
John Jayne

\$\$\$: NSF, DOE

